

Relationship among Nurses Role Overload, Burnout and Managerial Coping Strategies at Intensive Care Units

Fatma Rushdy Mohamed

Faculty of Nursing, Assiut University, 71515 Assiut, Egypt

Abstract

Work life takes up an important place in human life, and an employee faces many stimuli from internal and external environments and is affected by them in a positive or negative way.

Aim: was to investigate relationship among nurses role overload, burnout and managerial coping strategies at Intensive Care Units in Assiut University Hospital.

Design: A descriptive design was used.

Setting: Four Intensive Care Units in Assiut University Hospital.

Subjects: Included all nurses working in Casualty, General, & Postoperative Intensive Care Units and Coronary Care Unit in Assiut University Hospital.

Tools: Four tools which includes Socio-demographic data sheet, role overload questionnaire, managerial coping strategies & burnout questionnaires.

Results: Showed that role overload had highest mean score in General and Casualty ICUs. In addition, coping strategies highest mean scores of rational problem solving, resigned distancing, and seeking support/ventilation were in Casualty and post operative ICUs respectively except passive wishful thinking had highest mean scores in General and Casualty ICUs respectively. Also, burnout had highest mean scores in post operative and general ICU respectively.

Conclusions: Nurses employed at ICUs in Assiut University Hospital reported high burnout and use managerial coping strategies as rational problem solving, resigned distancing, seeking support/ventilation and passive wishful thinking to handle burnout and role overload except passive wishful thinking not used with role overload.

Background

Nurses having an important role in the health care system. they are considered to be members of a stressful job as a profession because they care for a stressful group comprising patients or those at health risk. The organizational structure of the health care system, improper working environment, critical patient care, working on a shift basis and the process of becoming a profession can affect the level of stress in nurses [1]. According to literature review the hardest hit by burnout are the service providers like nurses and counselors. When a professional is affected by burnout they feel drained and used up and have little desire to return to work the next day. This problem is specific to the work context. It is generally perceived that Intensive Care Nurses are particularly exposed to burnout since they literally deal with life and death situations most of the time. Therefore, it is suggested that nurses comprise the group that experiences the maximum stress, and burnout among health professionals. Burnout syndrome often occurs as a result of chronic work stress seen in Intensive Care Units [2].

The most widely referred definition of burnout among general occupations conceptualizes the phenomenon as "a syndrome of emotional exhaustion, cynicism and reduced professional efficacy". Researches suggest that burnout contributed to job stressors is associated with a negative outcome in both individuals and organizations. At an individual level, burnout has been associated with mental and physical health problems, for example, psychological distress, anxiety, depression, reduced self-esteem [3]. At an organization level, burnout is also consistently linked to negative attitudes towards work, which include low levels of motivation, job satisfaction and organizational commitment, but high levels of job uncertainty [4]. These in turn induce counter-productive work behavior, such as reduced productivity, absenteeism and staff turnover and hence lower the overall effectiveness of the organization itself [5].

Furthermore, some researches suggest that burnout is contagious, spreading to affect the colleagues of those who experience it and even resulting in negative spillover into one's home life [6]. From this standpoint, burnout is not only an individual well-being issue but could also influence socio-economic factors.

Coping behavior is a constantly changing cognitive and behavioral effort to manage specific external and/or internal demands that are appraised as taxing or exceeding the resources of the person. Coping behavior affects well-being and adaptation; In addition, coping behavior has also received attention from nursing researchers [7]. There is a growing understanding that coping strategies, which have been regarded as a psychological intervention in the stressor-strain relationship, play an important role in influencing the tangible and intangible outcomes of the stressor [8].

Coping strategies act as a variable that affects the direction and/or strength of the relation between an independent or predictor variable and a dependent or criterion variable²-it is reasonable to suggest that coping strategies may have a moderating effect on the relationship between the stressor and its consequential strain. Role overload may be regarded as a form of stressor, and burnout is one relevant stress reaction (i.e. strain), coping strategies are therefore likely to have a moderating effect on the relationship between role overload and

Corresponding Author: Dr. Fatma Rushdy Mohamed, Faculty of Nursing, Assiut University, 71515 Assiut, Egypt; E-mail: drfatmarushdy@yahoo.com

Citation: Mohamed FR (2016) Relationship among Nurses Role Overload, Burnout and Managerial Coping Strategies at Intensive Care Units. Int J Nurs Clin Pract 3: 180. doi: <http://dx.doi.org/10.15344/2394-4978/2016/180>

Copyright: © 2016 Mohamed. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

burnout. Therefore, given constant role overload levels, the presence of coping strategies weakens the stressor-strain relationship, reducing the levels of burnout in the individual [9].

Significance of the study

Nursing is a sensitive job which includes communicating with patients and caring after them; failing to identify and fighting the accompanying stress can have dire consequences. In such units as CCU and ICU the problems nurses encounter are more serious and they are subject to greater levels of stress. Obviously, the main reason for most physical and mental disorders is one's inability to deal with psychological pressure, which in turn affects one's reactions to stress [10]. The provision of Intensive Care (IC) can lead to a health care provider's physical, psychological and emotional exhaustion, which may develop into burnout. The researcher noticed that there was a limited specific study regarding this syndrome at Intensive Care Units (ICUs) in Assiut University Hospital. The findings will address some implication concerning the managerial coping strategies adopted by nurses at ICUs in the healthcare settings when dealing with job related burnout and role overload.

Aim of the study

The present study aims to investigate relationship among nurses role overload , burnout and managerial coping strategies (i.e. rational problem solving, resigned distancing, seeking support/ventilation and passive wishful thinking) at Intensive Care Units in Assiut University Hospital.

Research questions

- Is there a relationship between managerial coping strategies and nurses role overload?
- Is there a relationship between managerial coping strategies and nurses burnout?

Subjects and Methods

Study design

A descriptive design was used to carry out the present study.

Setting

The study was carried out at four Intensive Care Units in Assiut University Hospital which including the following: Causality Intensive Care Unit, General & Postoperative ICUs, and Coronary Care Unit (CCU).

Subjects

Includes all nurses working in aforementioned settings with a total number of 60 nurses classified as the following:

Tools of data collection

Four tools were used to collect data for this study as follows:

1st tool : Socio-demographic data sheet:

This part includes socio-demographic characteristics data as: age, sex , unit name, marital status, educational qualification, and years of experience.

Place of work	Number of Diploma degree Nurse	Number of Technical degree	Number of Bachelor degree Nurse Nurse
Causality Intensive Care Unit	-	6	15
General ICU	18	7	4
Postoperative ICU	10	10	-
Coronary Care Unit	16	10	4

2nd tool : Role overload questionnaire which developed by [11], it included five items. The Scoring system of this tool was five-point Likert-scale ranging from "strongly disagree" (1) to "strongly agree" (5).

3rd tool : Managerial coping strategies questionnaire which developed by [12] , it included twenty nine items classified into four factors. Rational problem solving, resigned distancing, seeking support/ventilation and passive wishful thinking. The Scoring system of the this study tool was four-point scale in the following sequence (0 = not used, 1= used somewhat, 2 = used quite a bit, 3 = used a great deal), and

4th tool: Burnout questionnaire which developed by [13], it included twenty two items. The Scoring system of this study tool was five-point Likert-scale ranging from "Not important" (1) to "More important" (5).

Their reliability was assessed in a pilot study by measuring their internal consistency using Cronbach's alpha coefficient method. This turned to be 0.88 for coping strategies, 0.90 for burnout, and 0.96 for role overload, thus indicating a high degree of reliability .The validity was measured by five experts in the field of nursing administration.

Pilot study

The pilot study also served to test the feasibility of the study and the clarity and practicability of the data collection tool. It was carried out on 20 nurses from different inpatient departments in Assiut University Hospital. The pilot study collected on April 2015. The subjects included in the pilot study were excluded. Data collected from the pilot study were reviewed and used in making the necessary modifications prior to the finalization of the data collection tool.

Fieldwork

An official permission was obtained from the hospital director, the nursing service director, and the head of each Intensive Care Unit before embarking on the study. The actual data collection was started in June 2015 and ended in July 2015. The researcher met the eligible nurses, explained to them the purpose of the study, and asked them for their oral consent to participate. Those who agreed to participate were given the tools and asked to fill them out and return them anonymously in the same setting or at most the next day.

Ethical considerations

The study protocol was approved by the pertinent authority. Participants' oral consent to participate was obtained after informing them about their rights to participate, refuse, or withdraw at any time. Total confidentiality of any obtained information was ensured. The study maneuver could not entail any harmful effects on participants.

Statistical analysis

Data entry and statistical analysis were done using SPSS 16.0 statistical software package. Data were presented using descriptive statistics. Pearson correlation analysis was used for assessment of the inter-relationships among quantitative variables, and Spearman rank correlation for ranked ones. Statistical significance was considered at p-value <0.05.

Table 1: Illustrated that more than three quarters of study subjects were aged less than 30 yrs and more than one quarter worked at Coronary Care Unit, 98% were females and half of them were single. While, 44% had a diploma degree and more than half of the study subject had less than 5 years of experience.

Table 2: Showed that role overload had highest mean score in general ICU followed by Casualty ICU (16.51±3.18 & 15.71±3.18). In addition, managerial coping strategies highest mean scores of rational problem solving, resigned distancing, and seeking support/ventilation were in Casualty ICU followed by post operative ICU (34.80±4.08& 33.50±7.68; 11.42±2.35 & 10.55± 0.32; and 15.23±3.40 & 14.50±3.76) respectively except passive wishful thinking had highest

mean scores in General ICU followed by Casualty ICU (12.68±3.08 & 12.28±3.63).Also, burnout had highest mean scores in post operative ICU followed by general ICU (66.30±14.54 & 63.17±14.72) respectively.

Table 3: Illustrates that burnout had positive relations with managerial coping strategies with a highly statistically significant differences (r = 0.49**). While, there were no relation between role overload & burnout (0.16) and role overload & coping strategies (0.13).

Concerning the relations of study variables with nurses' socio-demographic characteristics, Table 4) demonstrates that burnout had positive relations with role overload, rational problem solving, resigned distancing, seeking support/ventilation and passive wishful thinking coping strategies with a highly statistically significant differences (r = 0.49** , r = 0.34** , r =0.23* ,r = 0.32** , and r = 0.45**) respectively. While, role overload had weak negative statistically significant correlation with qualifications (r= -0.14). In addition, burnout had weak negative correlations with age, marital status, educational qualifications, and years of experience (r= 0.13, r= -0.02, r= -0.02& r= -0.14) respectively. Also, rational problem solving coping strategy had weak negative statistically significant correlations with age, marital status, and years of experience (r=-0.03, r= -0.23, & r= -0.06) respectively. Meanwhile, resigned distancing & seeking support/ventilation and passive wishful thinking coping strategies had weak negative statistically significant correlations with age and marital status(r= -0.09 , r= -0.18; r=-0.08, r= -0.19 & r= -0.12 , r= -0.26) respectively. There were positive relation among role overload and age, marital status, and years of experience (r = 0.04, r = 0.02 & r = 0.05) respectively. Educational qualifications had positive relations with the four managerial coping strategies rational problem solving, resigned distancing, seeking support/ventilation, and passive wishful thinking (r = 0.37** , r = 0.04, r = 0.20 &r = 0.07) respectively. Role overload had positive relations with rational problem solving, resigned distancing, and passive wishful thinking managerial coping strategies (r = 0.10, r = 0.14 & r = 0.07) respectively except seeking support/ventilation coping strategy had a negative relation with role overload (r = -0.01).

Discussion

The results of the present study showed that role overload had highest mean score in general ICU followed by Causality ICU. In addition, coping strategies highest mean scores of rational problem solving, resigned distancing, and seeking support/ventilation were in Causality ICU followed by post operative ICU respectively except passive wishful thinking had highest mean scores in General ICU followed by Causality ICU. Also, burnout had highest mean scores in post operative ICU followed by general ICU respectively (Table 2). These results indicated that nurses working in the intensive care unit, coronary care unit and emergency department experienced lower levels of personal accomplishment and high level of stress due to these nurses face more death and dying while exert more efforts. Nurses working in an intensive care unit rarely see patients returning home when hospital treatments are successful so nurses use coping strategies to overcome the effects of job stress.

These results is supported by a study on female hospital nurses, it was reported that affective-oriented coping was significantly associated with depressive symptoms, even after controlling for the effects of job stress, nursing-related work experience, and social support [14]. In addition, a study of hospital nurses in Australia and

	Frequency	Percent
Age (years):		
<30	79	79.0
30-35	14	14.0
>35		
7	7.0	
Mean±SD	25.28±0.58	
Department:		
Post operative ICU	20	20.0
General ICU	29	29.0
CCU	30	30.0
Causality ICU	21	21.0
Sex:		
Male	2	2.0
Female	98	98.0
Marital status:		
Single	50	50.0
Married	46	46.0
Divorced	3	3.0
Widow	1	1.0
Educational qualifications:		
Nursing Secondary school	44	44.0
Technical Institute of Nursing	33	33.0
Bachelor degree of Nursing	23	23.0
Experience years:		
<5	51	51.0
5-10	31	31.0
>10	18	18.0
Mean±SD	4.79 ±0.79	

Table 1: Socio-demographic characteristics of nurses in the study sample (n=100)

Study Variables	Mean + SD			
	Post operative ICU	General ICU	Coronary Care Unit	Causality ICU
Role overload	15.50±2.43	16.51±3.18	15.30±3.21	15.71±3.18
Managerial Coping strategies Rational problem solving	33.50±7.68	30.37±6.52	30.96±6.71	34.80±4.08
Resigned distancing	10.55± 0.32	10.27±2.76	10.43±2.67	11.42±2.35
Seeking support/ventilation	14.50±3.76	14.06±3.53	14.10±3.34	15.23±3.40
Passive wishful thinking	12.20±3.90	12.68±3.08	11.46±2.80	12.28±3.63
Burnout	66.30±14.54	63.17±14.72	60.90±12.89	57.38±15.23

Table 2: Mean scores of role overload, managerial coping strategies , and burnout among studied nurses at ICUs (n=100).

Variables	Pearson correlation coefficient		
	Scores		
	Role overload	Managerial Coping strategies	Burnout
Role overload		0.13	0.16
Managerial coping strategies	0.13		0.49**
Burnout	0.16	0.49**	

Table 3: Correlation of role overload, managerial coping strategies , and burnout among studied nurses at ICUs in Assiut University Hospital (n=100).

in the ICUs is hard than other general units because nurses in ICU provide intensive care for a long period (12 hours) shift and after that patients may died so these situation is stressful and lead to burnout. These results supported by [19] who mentioned that there are certain / inevitable stressful factors in nursing that can adversely affect nurses' minds and behaviors; it is important to find ways to deal with these factors before we can take measures to improve nurses' professional quality and teach them coping strategies. Nurses, when experiencing great stress, draw upon various coping strategies, and obviously, their manner of coping with stress and role overload influences the quality of their professional performance. In the same line the study results were in agreement with [20] who clarified that one of the important factors in avoiding burnout is applying stress coping

Characteristics	Spearman rank correlation coefficient					
	Scores of managerial coping strategies					
	Role overload	Rational problem solving	Resigned distancing	Seeking support/ventilation	Passive wishful thinking	Burnout
Age#	0.04	-0.03	-0.09	-0.08	-0.12	-0.13
Marital status (reference single)	0.02	-0.23*	-0.18	-0.19	-0.26**	-0.02
Qualification (reference: diploma)	-0.14	0.37**	0.04	0.20	0.07	-0.02
Experience years	0.05	-0.06	0.04	-0.08	-0.23*	-0.14
Role overload	-----	0.10	0.14	-0.01	0.07	0.49**
Burnout	0.49**	0.34**	0.23*	0.32**	0.45**	-----

Table (4):Correlation matrix for scores of role overload, coping strategies, burnout and socio-demographic characteristics of ICUs nurses(n=100)

(#) Pearson correlation coefficient

(*) Statistically significant at p<0.05

(**) Statistically significant at p<0.001

New Zealand found that type of coping strategy was significantly associated with physical and mental health, after controlling for the effects of job stress [15]. Moreover, [16] also reported that burnout among nurses is reduced when coping skills training was provided. A passive coping style, which is a type of emotion-focused coping, associates with poor health in both Norwegian and Dutch hospital nurses. Nurses with habitual passive coping may experience little control over work and low support within the nursing team [17]. Moreover, [3] concluded that positive stimuli in general make individuals happy whereas in the face of negative stimuli, stress can be generated. While, [18] found that burnout may develop in the individual as a result of a long duration of the state of stress.

On going the results of the current study demonstrates that burnout had positive relations with role overload and different types of managerial coping strategies with highly statistically significant differences (Table 3 & 4). These results might be attributed to the work

strategies. Moreover, [21] mentioned that two dimensions of coping are problem-focused coping and emotion-focused coping. Problem-focused coping consists of concrete attempts made to alter the stressfulevent while emotion-focused coping consists of attempts made to alter stressful feelings. People typically employ problem-focused coping strategies, purposively targeted at solving the problem at hand, when they perceive control over stressful events. Emotion-focused coping, aimed at minimizing negative emotions through seeking distraction and social support or by avoiding problems, predominates when people feel that the stressful event is something that must be endured.

As shown in (Table 4) burnout had weak negative correlations with age, marital status, educational qualifications, and years of experience. Also, rational problem solving coping strategy had weak negative statistically significant correlations with age, marital status, and years

of experience. Role overload had positive relations with rational problem solving, resigned distancing, and passive wishful thinking coping strategies respectively except seeking support/ventilation coping strategy had a negative relation with role overload. These results might be attributed to most of the studied nurses are young aged and had limited years of experience so they had not the ability to use the support /ventilation coping strategy and may be inability to communicate effectively which may reflected on their performance. This finding was inconsistent with [22] Foster, who found that older nurses in the United States experienced a lower level of burnout in emotional exhaustion, depersonalization, and reduced personal accomplishment than younger nurses.

In addition, A study in China found that Chinese nurses tended to cope actively with stress resulting from decreased personal achievement [23], but passively when the stress came from resource and environment problems, patient care and interaction issues, and from interpersonal relationships and management issues. Nurses apply different coping strategies, such as wishful thinking, planned problem-solving, to deal with different stressors, and apply avoidance strategies to cope with stress resulting from the shortage of resources, role overload and interpersonal relationships [23,24].

Conclusion

In the light of the study results, the following conclusions can be drawn:

Nurses employed at ICUs in Assiut University Hospital reported high burnout. They were more likely to use managerial coping strategies as rational problem solving, resigned distancing, seeking support/ventilation and passive wishful thinking to handle burnout. While, Role overload had positive relations with rational problem solving, resigned distancing, and passive wishful thinking managerial coping strategies except seeking support/ventilation coping strategy had a negative relation with nurses' role overload.

Recommendations

Based on the forgoing conclusions, the following recommendations are proposed:

- Recognizing the impacts of job-related stress and making use of effective coping methods play a vital role in reducing nurse's burnout.
- Developing and implementing a training program on managerial coping strategies to assist nursing staff in health service institution in Egypt to cope with burnout and role overload effectively.
- A change in leadership styles from the managerial level and reallocation of manpower may help reduce burnout and role overload.
- It is important for nurse managers to recognize passive coping, because this type of coping associates with poor health
- Hospital decision-makers need to consider arranging for the nurses to work in other units for periods of time during the year to buffer their negative experience at work.
- Further researches should be done to investigate relations among job satisfaction, job stress, personality traits and burnout among ICU staff.

Competing Interest

The authors declare that they have no competing interests.

References

1. Tuna R, Bayka U (2014) The relationship between job stress and burnout levels of oncology nurses. *Asia-Pacific Journal of Oncology Nursing* 1: 33-39.
2. Chitura D, Chitura M (2014) Burnout syndrome in intensive care unit nurses in Zimbabwe. *European Scientific Journal* 436- 457.
3. Maslach C, Schaufeli W, Leiter M (2001) Job burnout. *Annual Review of Psychology* 52: 397-422.
4. Yip B, Rowlinson S (2006) Job burnout among construction professionals in Hong Kong. *Journal of the Hong Kong Institute of Construction Managers* 11-14.
5. Wright T, Bonett D (1997) The contribution of burnout to work performance. *Journal of Organizational Behavior* 18: 491-499.
6. Westman M, Eden D (1997) Effects of respite from work on burnout: vacation relief and fade out. *J Appl Psychol* 82: 516-527.
7. Kato T (2014) Coping with interpersonal stress and psychological distress at work: comparison of hospital nursing staff and salespeople. *Psychol Res Behav Manag* 7: 31-36.
8. Tidd S, Friedman R (2002) Conflict style and coping with role conflict: an extension of the uncertainty model of work stress. *International Journal of Conflict Management* 13: 236-257.
9. Yip Y, Rowlinson S, Siu O (2008) Coping strategies as moderators in the relationship between role overload and burnout. *Construction Management and Economics* 26: 871-882.
10. Akouchekian S, Roohafza H, Hasanzadeh A, Sharifi HM (2009) Relation between Social Support and Coping with Stress in Nurses in Psychiatric Ward. *Journal of Guilan University of Medical Sciences* 18: 41-46.
11. Harris M, Bladen A (1994) Wording effects in the measurement of role conflict and role ambiguity: a multi-trait-multi-method analysis. *Journal of Management* 20: 887-901.
12. Chan D (1994) The Chinese ways of coping questionnaire: assessing coping in secondary teachers and students in Hong Kong. *Psychological Assessment* 6: 108-116.
13. Maslach C, Jackson S, Leiter M (1996) *Maslach Burnout Inventory Manual*, Palo Alto, CA: Consulting Psychologists Press. 3rd edn.
14. Lin H, Probst J, Hsu Y (2010) Depression among female psychiatric nurses in southern Taiwan: main and moderating effects of job stress, coping behavior and social support. *J Clin Nurs* 19: 2342-2354.
15. Chang E, Bidewell J, Huntington A (2007) A survey of role stress, coping and health in Australian and New Zealand hospital nurses. *Int J Nurs Stud* 44: 1354-1362.
16. Gunusen N, Ustun B (2010) An RCT of coping and support groups to reduce burnout among nurses. *Int Nurs Rev* 57: 485-492.
17. Schreuder J, Roelen C, Groothoff J, Van der Klink J, Mageroy N, et al. (2012) Coping styles relate to health and work environment of Norwegian and Dutch hospital nurses: A comparative study. *Nursing outlook* 60: 37-43.
18. Spooner-Lane R, Patton W (2007) Determinants of burnout among public hospital nurses. *Aust J Adv Nurs* 25: 8-16.
19. Lashonda L (2004) Factors that most influence job satisfaction among cardiac nurses in an acute setting. [Thesis]. USA. Marshall University.
20. GrusiFarshi M, Moslemi M (2005) Investigate the relationship between burnout and coping techniques Nurses Kurdistan province. *Quarterly Journal of Psychological Studies* 1: 5-19.
21. LeSergenta C, Haney C (2005) Rural hospital nurse's stressors and coping strategies: a survey. *Int J Nurs Stud* 42: 315-324.
22. Foster E (2003) Stress, burnout and coping strategies of nurses practicing in two state public psychiatric hospitals in the New York metropolitan area. Waldon University (Dissertation).

23. Xie W, Zhang Z, Zheng Z, Liang B (2005) Relationship among job burnout, job stressors, and coping ways of nurses. Chinese Journal of Behavioral Medical Science 14: 462-463.
24. Bao M, Liu Y, Wu J (2011) Study on job stressors and coping ways of Bachelor nurses (Article in Chinese). Chinese Journal of Disease Control and Prevention 16: 937-939.